

(<https://www.AnesthesiologyNews.com>)



(https://www.facebook.com/dialog/feed?app_id=1526701630986537&display=popup&caption=Unplanned or Accidental Extubation in the Perioperative Environment&link=https://www.anesthesiologynews.com/Review-Articles/Article/08-19/Unplanned-or-Accidental-Extubation-in-the-Perioperative-Environment/55658&redirect_uri=https://www.anesthesiologynews.com/Review-Articles/Article/08-19/Unplanned-or-Accidental-Extubation-in-the-Perioperative-Environment/55658)



(<https://twitter.com/intent/tweet?text=Unplanned or Accidental Extubation in the Perioperative Environment%0D%0A%0D%0Ahttps://www.anesthesiologynews.com/Review-Articles/Article/08-19/Unplanned-or-Accidental-Extubation-in-the-Perioperative-Environment/55658>)



([https://www.linkedin.com/shareArticle?mini=true&url=https://www.anesthesiologynews.com/Review-Articles/Article/08-19/Unplanned-or-Accidental-Extubation-in-the-Perioperative-Environment/55658&title=Anesthesiology News - Unplanned or Accidental Extubation in the Perioperative Environment&summary=More than 120,000 incidents of unplanned or uncontrolled extubation occur yearly in the perioperative environment and ICU, resulting in significant complications, increased costs and increased hospital lengths of stay. \)](https://www.linkedin.com/shareArticle?mini=true&url=https://www.anesthesiologynews.com/Review-Articles/Article/08-19/Unplanned-or-Accidental-Extubation-in-the-Perioperative-Environment/55658&title=Anesthesiology News - Unplanned or Accidental Extubation in the Perioperative Environment&summary=More than 120,000 incidents of unplanned or uncontrolled extubation occur yearly in the perioperative environment and ICU, resulting in significant complications, increased costs and increased hospital lengths of stay.)))

More than 120,000 incidents of unplanned or uncontrolled extubation occur yearly in the perioperative environment and ICU, resulting in significant complications, increased costs and increased hospital lengths of stay.)

Review Articles (/Section/Review-Articles/161)

AUGUST 27, 2019

Unplanned or Accidental Extubation in the Perioperative Environment


Lauren Berkow, MD, FASA

*President, Society for Airway Management
Professor of Anesthesiology
University of Florida College of Medicine
Gainesville, Fla.*

Dr. Berkow is a member of the Teleflex Medical Advisory Board and the Masimo Scientific Advisory Board.

Download the printer-friendly PDF version here. (http://cms.mcmahonmed.com//download/Unplanned_ANSE0819_WM.pdf)

More than 120,000 incidents of unplanned or uncontrolled extubation occur yearly in the perioperative environment and ICU, resulting in significant complications, increased costs and increased hospital lengths of stay.



(/images/2019/ANSE0819_0071_a_600.jpg)

Introduction

Endotracheal intubation and extubation are procedures routinely performed by anesthesiologists and intensive care physicians. The majority of the time, extubation is a planned event. Even when planned, the rate of complications related to extubation have been reported in the literature to be as high as 12%.^{1,2} Incidents of unplanned extubation can occur in the OR setting, the ICU, the emergency department, or any specialty department (medical imaging, endoscopy suite, interventional cardiology) in which an intubated patient may need to be cared for. "Unplanned extubation" is defined as the unintentional and uncontrolled removal of the endotracheal tube (ETT) either due to actions of the patient or during nursing care or movement of the patient.³⁻⁵ Extensive publications exist in the literature devoted to prediction and management of difficult intubation, yet both planned and unplanned extubations have been much less widely studied, especially outside the OR environment where airway complications are higher.⁶⁻⁷

ADVERTISEMENT

Incidence and Risk Factors

In the literature, the reported incidence of unplanned extubation ranges from 0.5% to 35.8% in adults.^{4,8-13} The majority of studies have been conducted in the ICU setting because data from outside the ICU are negligible. In the neonatal ICU (NICU), unplanned extubation has been reported as the fourth most common adverse event, and rates have been reported to be as high as 80.8%.¹⁴⁻¹⁷

On average, in the United States, 1.65 million adult patients are mechanically ventilated each year in the ICU.^{18,19} Based on a median unplanned extubation rate of 7.3%, more than 120,000 incidents of unplanned or uncontrolled extubation occur yearly in the ICU.⁴ In the NICU, 445,000 patients are mechanically ventilated. Based on the median unplanned extubation rate in the NICU of 18.2%, more than 80,000 neonates experience an unplanned extubation yearly.¹⁶ Outside the ICU, the rate of occurrence is unknown, as published studies are scant.

Why might an unplanned extubation occur in the OR? Occasionally, the patient self-extubates upon emergence. Typically, these patients do not need reintubation, but vocal cord injury may occur if the ETT cuff was not deflated. More concerning are the rare events of unplanned or accidental extubation during the operative procedure. This could occur during lateral or prone positioning, or during operative procedures in or proximate to the airway. Several case reports have described accidental extubation while the patient was in the prone position for spine surgery.¹⁸⁻²⁰ In one case, extubation occurred during a "wake-up" test to assess neurologic function and was successfully managed with a supraglottic airway device (SAD).²⁰ In another case, extubation occurred while the patient was prone with the head flexed and secured with surgical pins, requiring fiber-optic intubation (FOI).²¹

Procedures in which the bed is turned 180 degrees away from the anesthesia machine also present a higher risk, since the ETT cannot be directly visualized or monitored during the case. Spond et al reported a case of a patient with Klippel-Feil syndrome undergoing prone cervical fusion with the bed turned 180 degrees away from the anesthesia machine, who was accidentally extubated during the procedure.²² In this case report, with a known difficult airway, the patient was flipped supine to perform FOI.

Why do unplanned extubations occur in the intensive care setting? The most common cause in adults is self-extubation by the patient (62%-96% of incidents).^{23,24} Other common causes include accidental extubation during suctioning, manipulation of the ETT, or moving the patient.⁴ Unplanned extubation can occur during turning or repositioning of the patient. When intubated patients need to travel from the ICU for a diagnostic or an interventional procedure, they are moved several times (such as for a CT or MRI scan), providing multiple opportunities for dislodgement or removal of the ETT.

Several risk factors can increase the likelihood of extubation by accident or patient action (Table 1).^{4,8,11,25-29} Inadequate securement of the tube can increase the risk for removal or dislodgement. Lack of physical restraints, inadequate patient sedation, or patient agitation or restlessness in the setting of an inadequately secured ETT also increase the risk for tube dislodgement or removal.^{4,8,11} Other risk factors that have been linked to unplanned extubation in the intensive care setting include emergency surgery, presence of nosocomial infection, delirium or confusion, congestive heart failure, and lack of a clear plan for extubation.²⁶⁻²⁸

Table 1. Risk Factors for Unplanned Extubation^{4,8,11,18-21}

Inadequate securement of the endotracheal tube
Inadequate sedation
Lack of physical restraints
Patient restlessness or agitation
Unclear extubation plan

Absence of clear extubation or weaning policies and procedures also has been associated with an increased risk for unplanned extubations.^{4,8,25} Other human factors, such as fatigue, inadequate staffing patterns in the ICU, and level of nursing experience have been linked to unplanned or accidental extubation.^{4,25} Nursing workloads and higher nurse-to-patient ratios also have been shown to contribute to unplanned extubation.^{24,29} Not surprisingly, the incidence of unplanned extubation is higher during evening and night shifts.³⁰

Danielis et al interviewed critical care nurses about precipitating factors for unplanned extubation.²⁹ They reported lack of communication between health care providers, a chaotic environment, poor nurse-to-patient ratios, and barriers to the ability to directly observe the patient as significant factors.

Complications

In the OR, a significant number of reported airway-related complications occur during extubation. According to the American Society of Anesthesiologists' closed claims database, 12% of all airway claims were associated with extubation.^{1,31} The 4th National Audit Project in the United Kingdom, which collected airway-related claims reported by anesthesiologists, found that one-third of claims were due to respiratory complications during emergence and extubation.³² These were complications related to planned extubation in the OR setting, under presumed controlled conditions. Complication rates related to unplanned extubation are even higher.

Unplanned extubation can lead to a large variety of complications (Table 2).^{24,33,34} Unplanned removal of an ETT with the tracheal cuff still inflated can potentially cause injury to the vocal cords. If an intubated patient has a large secretion burden, aspiration of these secretions after an unplanned extubation can lead to aspiration pneumonia. If inadequate ventilation occurs after unplanned extubation, hypoxemia and potentially hemodynamic instability, hypotension, brain damage, cardiac arrest, and even death may result.

Table 2. Complications Associated With Unplanned Extubation^{22,29-30}

Clinical Complications	Other Complications
Aspiration pneumonia	Increased hospital costs
Brain damage	Increased hospital LOS
Cardiac arrest	Increased ICU mortality rates
Death	Prolonged ICU LOS
Hemodynamic instability	
Hypoxemia	
Respiratory failure	
Vocal cord injury	

LOS, length of stay

Reintubation after unplanned extubation, especially in the presence of airway edema, can be very challenging, and can result in the previously mentioned complications. The reintubation rate after unplanned extubation varies in the literature, but has been reported to be as high as 89%.^{4,25,33,35} A study by Mort on unplanned extubations in the ICU found that 89% of patients required reintubation within two hours, and 66% needed to be reintubated within 30 minutes of accidental extubation.³⁶ Patients with accidental extubation, as opposed to self-extubation, have been reported to have higher reintubation rates as well as poorer prognoses.⁷³ A study by de Lassence et al found an overall reintubation rate of 77%, with the majority of patients after accidental extubation requiring reintubation, while 37% of self-extubated patients were not reintubated.²⁴

Ventilator-associated pneumonia (VAP) has been clearly linked to prolonged mechanical ventilation and longer ICU and hospital lengths of stay, independent of the cause.^{38,39} Unplanned extubation is a risk factor and has been associated with increased rates of ventilator-associated pneumonia.⁴⁰ Unplanned extubation, due to the subsequent complications, has been associated with increased ICU and hospital length of stay (LOS), as well as increased mortality.^{24,41-43} Other studies, however, found no difference in ICU LOS or mortality rates compared with intubated patients who did not experience unplanned extubation.^{8,35} The study by de Lassence et al examined LOS and found that in adult patients with

unplanned extubation, both ICU LOS (18 vs. planned days) and total LOS (30 vs. 18 days) were prolonged. A similar result was found in pediatric patients with unplanned extubation.^{44,45} Veldman reported an increased NICU LOS of 51 versus nine days.⁴⁴

More severe complications also have been associated with unplanned extubation. A study by Klugman found that 20% of unplanned extubations were followed by cardiovascular collapse, with neonates being at higher risk.⁴⁶ Multiple studies have reported increases in mortality rate and poorer prognosis in patients with unplanned extubation.^{35,42,47}

Costs

Increases in hospital and ICU LOS as well as in complications all lead to increased hospital costs that can be directly attributed to unplanned extubation.⁴⁸⁻⁵⁰ Mechanical ventilation in the ICU, independent of any complications associated with it, increases costs. Dasta et al assessed costs in the ICU setting and found that mechanical ventilation increased costs by an average of \$1,522 per day.⁴⁹ With the average LOS of a patient experiencing an unplanned extubation being 30 days, the total increased cost of an unplanned extubation is \$40,992. The complications associated with unplanned extubation in this patient population further increase hospital costs. Roddy et al calculated the costs of unplanned extubation in the pediatric ICU setting, factoring in increased LOS and rate of nosocomial infection, and found that unplanned extubation increased hospital costs by over \$36,000 per incident.⁵⁰

Taking into account mortality, increased rates of VAP and increased LOS, the overall cost burden in the United States of unplanned extubation in adult ICUs totals almost \$5 billion (Table 3).^{24,48,50} In the NICU, the more than 80,000 unplanned extubations yearly costs an additional \$2.9 billion.⁵⁰

Table 3. Average Yearly Costs Associated With Unplanned Extubation in the ICU^{24,48,50}

Day	Non-UE (\$)	UE (\$)
1	10,794	10,794
2	4,796	4,796
3	3,968	3,968
4	3,968	3,968
5	3,968	3,968
6	3,968	3,968
7	3,968	3,968
8	3,968	3,968
9	3,968	3,968
10	1,760	3,968
11	1,760	3,968
12	1,760	3,968
13	1,760	3,968
14	1,760	3,968
15	1,760	3,968
16	1,760	3,968
17	1,760	3,968
18	1,760	3,968
19		1,760
20		1,760
21		1,760
22		1,760
23		1,760
24		1,760
25		1,760
26		1,760
27		1,760
28		1,760

30	1,760
59,206	100,198
Average cost of UE	40,992
Average yearly No. of UEs	121,000
Yearly cost of UEs	4,960,032,000
<input type="checkbox"/> ICU Days on MV	
<input type="checkbox"/> ICU Days off MV	
<input type="checkbox"/> Non-ICU Days	
MV, mechanical ventilation; UE, unplanned extubation	

Prevention

Surprisingly, this important problem remains underrecognized. Many hospitals still do not track rates of unplanned extubation. A major first step in prevention is to increase awareness of the problem. Several strategies can be implemented to reduce the risk for unplanned extubation.

Increased Awareness of Risk Factors

The first step to increase awareness is education of the staff providing care to intubated patients about risk factors for unplanned extubation and how to monitor patients at risk. Vats et al created an airway risk assessment scoring tool to identify and stratify pediatric patients at risk for unplanned extubation in the ICU.⁵¹ Patients were assigned points for several risk factors:

- anatomic risk;
- secretions;
- agitation;
- prone position;
- history of unplanned extubation;
- need for multiple procedures or transports; and
- frequent need for retaping of the ETT.

A score of 5 or greater was considered high risk, and the investigators found that the scoring tool correlated well with the incidence of unplanned extubation.

Several institutions have tackled prevention of unplanned extubation as a quality improvement initiative, incorporating multidisciplinary interventions to affect outcomes.^{24,26,52,53} Chao et al reported a significant decrease in unplanned extubation rates, from 3.19 to 0.95 per 100 patients, using a multidisciplinary strategy that focused on these areas: standardization of procedures, improving communication, revising sedation and weaning protocols, changing restraint strategies and securement methods, using team resource management and a no-blame culture, and creation of a task force for identifying high-risk patients.⁵² Quality improvement initiatives have also been successful in reducing unplanned extubation rates in pediatric ICUs.^{54,55}

Strategies to Reduce the Incidence of Unplanned Extubation

Individual strategies also can reduce the incidence of unplanned extubation. Standardization of protocols as well as the creation of bundles and checklists in the ICU setting have been shown to reduce VAP and improve outcomes.^{56,57} Standardization of restraint and sedation protocols could have a similar benefit for unplanned extubation.⁵²

Improving and optimizing securement of the ETT in the ICU can prevent unplanned extubation. Many different methods and securement devices exist to maintain an indwelling ETT. Although no single method or device has been proven superior, there are several attributes that an optimal securement method should have (Table 4).⁵⁸⁻⁶⁰

<p>Table 4. Attributes of an Optimal ETT Securement Device⁵⁷⁻⁵⁹</p> <p>Adequate stabilization against external forces that may dislodge tube</p> <p>Allows tube movement for oral care</p> <p>Avoids adhesives that could irritate skin</p> <p>Cost-effective</p>

Ease of placement, use, and maintenance
Enhances patient comfort and minimizes skin pressure
Facilitates suctioning but is not compromised by secretions
Prevents tube movement
Requires infrequent adjustment or change
ETT, endotracheal tube

Addressing Human and Environmental Factors

Human factors such as staffing ratios, interprofessional communication and teamwork should be addressed and optimized to reduce risk in both the OR and ICU settings. The simulation environment can be very useful for practicing and teaching teamwork and communication.^{61,62} Danielis et al surveyed nurses in the ICU and identified several organizational, environmental and nursing care factors that played a role in unplanned extubations and could be modified.²⁹ These factors included communication failures, environmental chaos and barriers to direct surveillance of the patient, and poor nurse-to-patient ratios.²⁹

Operating Room Strategies

In the OR, a briefing or time-out can be used to discuss extubation risk prevention and strategies for management in high-risk cases, similar to what is recommended for cases at risk for an airway fire.²² Many cases at risk for an airway fire may also be at risk for accidental extubation during the procedure. Careful securement of the ETT for cases in which the airway is inaccessible during the procedure (surgery on the head and neck, prone or lateral procedures, procedures where the bed is turned 90 or 180 degrees away from the anesthesia machine) can potentially prevent extubation. In cases with high risk, immediate availability of alternate airway devices such as video laryngoscopes, flexible bronchoscopes and SADs is recommended for emergent airway management.²⁰⁻²²

Prevention of Reintubation After Unplanned Extubation

There are several strategies to maximize oxygenation and ventilation after unplanned extubation that could reduce the need for reintubation, or at least result in a less urgent need for reintubation. Newer methods of high-flow oxygenation via the nasal route that also provide positive pressure ventilation may provide a bridge after unplanned extubation.^{63,64} Use of continuous positive airway pressure after extubation, both planned and unplanned, also may be useful, especially in obese patients or individuals with obstructive sleep apnea.^{65,66} Lin et al found that the use of noninvasive positive pressure ventilation significantly reduced the reintubation rate after unplanned extubation.³⁵

Conclusion and Future Directions

Unplanned extubation is a common and costly problem in the perioperative and intensive care environments, with a large impact on outcomes and hospital costs, yet it remains an underrecognized problem. Increased awareness and prevention are critical. Better tracking and the implementation of quality improvement initiatives can potentially address the problem. Prevention requires commitment not only from clinical care providers but also from leadership, to implement strategies and protocols to standardize care. Rates of unplanned extubation should be identified and tracked, ideally within an electronic health record system.

The Society for Airway Management has collaborated with many medical societies and safety organizations to address the issue of unplanned extubation (Table 5). This collaborative is focused on increasing awareness of the magnitude of the problem as well as improved tracking of events. The collaborative also is developing checklists, core data sets for tracking and Actionable Patient Safety Solutions in conjunction with the Patient Safety Movement Foundation.⁶⁷ This information can be downloaded at www.airwaysafety movement.org (<http://www.airwaysafety movement.org>) or www.patientsafety movement.org (<http://www.patientsafety movement.org>).

Table 5. Society for Airway Management Collaborative
Medical Professional Societies
American Academy of Pediatrics
American Association of Nurse Anesthetists
American Association of Respiratory Care
American College of Emergency Physicians
American Society of Anesthesiologists
Association of Air Medical Systems

National Association of EMTs
National Association of Neonatal Nurses
Society for Airway Management
Society for Pediatric Anesthesia
Society of Critical Care Medicine
Patient Safety Organizations
Airway Safety Movement
Anesthesia Patient Safety Foundation
Children's Hospitals' Solutions for Patient Safety
Do It For Drew Foundation
Emergency Medicine Patient Safety Foundation
Patient Safety Movement Foundation
Quality Improvement Organizations
IMPAQ/CMS Strategic Innovation Engine

References

- Asai T, Koga K, Vaughan RS. Respiratory complications associated with tracheal intubation and extubation. *Br J Anaesth*. 1998;80(6):767-775.
- Domino KB, Posner KL, Caplan RA, et al. Airway injury during anesthesia: a closed claims analysis. *Anesthesiology*. 1999;91(6):1703-1711.
- Ismaeil MF, El-Shahat HM, El-Gammal MS, et al. Unplanned versus planned extubation in respiratory intensive care unit, predictors of outcome. *Egyptian J Chest Dis Tuberc*. 2014;63(1):219-231.
- da Silva PS, Fonseca MC. Unplanned endotracheal extubations in the intensive care unit: systematic review, critical appraisal, and evidence-based recommendations. *Anesth Analg*. 2012;114(5):1003-1014.
- Bouza C, Garcia E, Diaz M, et al. Unplanned extubation in orally intubated medical patients in the intensive care unit: a prospective cohort study. *Heart Lung*. 2007;36(4):270-276.
- Cook T, Behringer EC, Bengler J. Airway management outside the operating room: hazardous and incompletely studied. *Curr Opin Anaesthesiol*. 2012;25(4):461-469.
- Asai T. Airway management inside and outside operating rooms—circumstances are quite different. *Br J Anaesth*. 2018;120(2):207-209.
- de Groot RI, Dekkers OM, Herold IH, et al. Risk factors and outcomes after unplanned extubations on the ICU: a case-control study. *Crit Care*. 2011;15(1):R19.
- de Lassence A, Alberti C, Azoulay E, et al. Impact of unplanned extubation and reintubation after weaning on nosocomial pneumonia risk in the intensive care unit: a prospective multicenter study. *Anesthesiology*. 2002;97(1):148-156.
- Phoa LL, Pek WY, Syap W, et al. Unplanned extubation: a local experience. *Singapore Med J*. 2002;43(10):504-508.
- McNett M, Kerber K. Unplanned extubations in the ICU: Risk factors and strategies for reducing adverse events. *J Clin Outcomes Manag*. 2015;22(7):303-311.
- Kapadia FN, Bajaj KB, Raju KV. Airway accidents in intubated intensive care unit patients: an epidemiological study. *Crit Care Med*. 2000;28(3):659-664.
- Balon JA. Common factors of spontaneous self-extubation in a critical care setting. *Int J Trauma Nurs*. 2001;7(3):93-99.
- Meyers JM, Pinheiro J, Nelson MU. Unplanned extubation in NICU patients: are we speaking the same language? *J Perinatol*. 2015;35(9):676-677.
- Barber JA. Unplanned extubation in the NICU. *J Obstet Gynecol Neonatal Nurs*. 2013;42(2):233-238.
- Silva PS, Reis ME, Aguiar VE, et al. Unplanned extubation in the neonatal ICU: a systematic review, critical appraisal, and evidence-based recommendations. *Respir Care*. 2013;58(7):1237-1245.
- Franck LS, Vaughan B, Wallace J. Extubation and reintubation in the NICU: identifying opportunities to improve care. *Pediatr Nurs*. 1992;18(3):267-270.
- Society of Critical Care Medicine. Critical care statistics. www.sccm.org/Communications/Critical-Care-Statistics (<http://www.sccm.org/Communications/Critical-Care-Statistics>). Accessed April 10, 2019.
- Wunsch H, Wagner J, Herlim M, et al. ICU occupancy and mechanical ventilator use in the United States. *Crit Care Med*. 2013;41(12):2712-2719.
- Rapheal J, Rosenthal-Ganon T, Gozal Y. Emergency airway management with a laryngeal mask airway in a patient placed in the prone position. *J Clin Anesth*. 2004;16(7):560-561.
- Hung MH, Fan SZ, Lin CP, et al. Emergency airway management with fiberoptic intubation in the prone position with a fixed flexed neck. *Anesth Analg*. 2008;107(5):1704-1706.
- Spond M, Burns T, Rosenbaum T, et al. Crisis management of accidental extubation in a prone-positioned patient with Klippel-Feil syndrome. *A & A Case Rep*. 2016;6(12):383-386.
- Kiekkas P, Aretha D, Panteli E, et al. Unplanned extubation in critically ill adults: clinical review. *Nurs Crit Care*. 2013;18(3):123-134.
- de Lassence A, Alberti C, Azoulay E, et al. Impact of unplanned extubation and reintubation after weaning on nosocomial pneumonia risk in the intensive care unit. A prospective multicenter study. *Anesthesiology*. 2002;97(1):148-156.
- Cosentino C, Fama M, Foà C, et al. Unplanned extubations in intensive care unit: evidences for risk factors. A literature review. *Acta Biomed*. 2017;88(5S):55-65.
- Ai ZP, Gao XL, Zhao XL. Factors associated with unplanned extubation in the intensive care unit for adult patients: A systematic review and meta-analysis. *Intensive Crit Care Nurs*. 2018;47:62-68.
- Aydogan S, Kaya N. The assessment of the risk of unplanned extubation in an adult intensive care unit. *Dimens Crit Care Nurs*. 2017;36(1):14-21.
- Piriyapatsom A, Chittawatanarat K, Kongsayreepong S, et al. Incidence and risk factors of unplanned extubation in critically ill surgical patients: the multi-center Thai University-based surgical intensive care units study (THAI-SICU Study). *J Med Assoc Thai*. 2016;99(suppl 6):S153-S162.
- Danielis M, Chiaruttini S, Palese A. Unplanned extubations in the intensive care unit: findings from a critical incident technique. *Intensive Crit Care Nurs*. 2018;47:69-77.
- Kwon E, Choi K. Case-control study on risk factors of unplanned extubation based on patient safety model in critically ill patients with mechanical ventilation. *Asian Nurs Res*. 2017;11(1):74-78.
- Metzner J, Posner KL, Lam MS, et al. Closed claims' analysis. *Best Pract Res Clin Anaesthesiol*. 2011;25(2):263-276.
- Cook TM, Woodall N, Frerk C. Major complications of airway management in the UK: results of the 4th National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 1: anaesthesia. *Br J Anaesth*. 2011;106(5):617-631.
- Chao CM, Sung MI, Cheng KC, et al. Prognostic factors and outcomes of unplanned extubation. *Sci Rep*. 2017;7(1):8636.
- Kapadia F. Effect of unplanned extubation on outcome of mechanical ventilation. *Am J Respir Crit Care Med*. 2001;163(7):1755-1756.
- Lin PH, Chen CF, Chiu HW, et al. Outcomes of unplanned extubation in ordinary ward are similar to those in the intensive care unit. A STROBE-compliant case-control study. *Medicine*. 2019;98(11):e14841.
- <https://www.anesthesiologynews.com/Review-Articles/Article/08-19/Unplanned-or-Accidental-Extubation-in-the-Perioperative-Environment/55658>


37. Bhattacharya P, Chakraborty A, Agarwal P. Comparison of outcome of self-extubation and accidental extubation in ICU. *Indian J Crit Care Med*. 2007;11(3):105-108.
38. Rosenthal VD, Udwardia FE, Muñoz HJ, et al. Time-dependent analysis of extra length of stay and mortality due to ventilator-associated pneumonia in intensive-care units of ten limited-resources countries: findings of the International Nosocomial Infection Control Consortium (INICC). *Epidemiol Infect*. 2011;139(11):1757-1763.
39. Karvouniaris M, Makris D, Manoulakas E, et al. Ventilator-associated tracheobronchitis increases the length of intensive care unit stay. *Infect Control Hosp Epidemiol*. 2013;34(8):800-808.
40. Gao F, Yang LH, He HR, et al. The effect of reintubation on ventilator-associated pneumonia and mortality among mechanically ventilated patients with intubation: a systematic review and meta-analysis. *Heart Lung*. 2016;45(4):363-371.
41. Chuang ML, Lee CY, Chen YF, et al. Revisiting unplanned endotracheal extubation and disease severity in intensive care units. *PLoS One*. 2015;10(10):e0139864.
42. Epstein SK, Nevins ML, Chung J. Effect of unplanned extubation on outcome of mechanical ventilation. *Am J Respir Crit Care Med*. 2000;161(6):1912-1916.
43. Elmetwally R, Elshafey M, Zeidan A, et al. Predictors and outcome of unplanned extubation in COPD patients. *Egypt J Anaesth*. 2010;26(1):1-11.
44. Veldman A, Trautschold T, Weiss K, et al. Characteristics and outcome of unplanned extubation in ventilated preterm and term newborns on a neonatal intensive care unit. *Paediatr Anaesth*. 2006;16(9):968-973.
45. Kanthimathinathan HK, Durward A, Nyman A, et al. Unplanned extubation in a paediatric intensive care unit: prospective cohort study. *Intensive Care Med*. 2015;41(7):1299-1306.
46. Klugman D, Berger JT, Spaeder MC, et al. Acute harm: Unplanned extubations and cardiopulmonary resuscitation in children and neonates. *Intensive Care Med*. 2013;39(7):1333-1334.
47. Lee JH, Lee HC, Jeon YT, et al. Clinical outcomes after unplanned extubation in a surgical intensive care population. *World J Surg*. 2014;38(1):203-210.
48. Dasta JF, McLaughlin TP, Mody et al. Daily cost of an intensive care unit day: The contribution of mechanical ventilation. *Crit Care Med*. 2005;33(6):1266-1271.
49. Needham DM, Pronovost PJ. The importance of understanding the costs of critical care and mechanical ventilation. *Crit Care Med*. 2005;33(6):1434-1435.
50. Roddy DJ, Spaeder MC, Pastor W, et al. Unplanned extubations in children: impact on hospital cost and length of stay. *Pediatr Crit Care Med*. 2015;16(6):572-575.
51. Vats A, Hopkins C, Hatfield K, et al. An airway risk assessment score for unplanned extubation in intensive care pediatric patients. *Pediatr Crit Care Med*. 2017;18(7):661-666.
52. Chao CM, Lai CC, Chan KS, et al. Multidisciplinary interventions and continuous quality improvement to reduce unplanned extubation in adult intensive care units. A 15-year experience. *Medicine*. 2017;96(27):e6877.
53. Chiang AA, Lee KC, Lee JC, et al. Effectiveness of a continuous quality improvement program aiming to reduce unplanned extubation: a prospective study. *Intensive Care Med*. 1996;22(11):1269-1271.
54. Tripathi S, Nunez DJ, Katyal C, et al. Plan to have no unplanned: a collaborative, hospital-based quality-improvement project to reduce the rate of unplanned extubations in the pediatric ICU. *Respir Care*. 2015;60(8):1105-1112.
55. Fontanez-Nieves TD, Frost M, Anday E, et al. Prevention of unplanned extubations in neonates through process standardization. *J Perinatol*. 2016;36(6):469-473.
56. Scales DC, Dainty K, Hales B, et al. A multifaceted intervention for quality improvement in a network of intensive care units: a cluster randomized trial. *JAMA*. 2011;305(4):363-372.
57. Bouadma L, Deslandes E, Lolom I, et al. Long-term impact of a multifaceted prevention program on ventilator-associated pneumonia in a medical intensive care unit. *Clin Infect Dis*. 2010;51(10):1115-1122.
58. Fisher DF, Chenelle CT, Marchese AD, et al. Comparison of commercial and noncommercial endotracheal tube-securing devices. *Respir Care*. 2014;59(9):1315-1323.
59. Gardner A, Hughes D, Cook R, et al. Best practice in stabilisation of oral endotracheal tubes: a systematic review. *Aust Crit Care*. 2005;18(4):158, 160-165.
60. Mehta NM, Sharma S, Laussen PC. Unplanned extubation: securing the tool of our trade. *Intensive Care Med*. 2015;41(11):1983-1985.
61. Umoren RA, Poore JA, Sweigart L, et al. TeamSTEPPS virtual teams: interactive virtual team training and practice for health professional learners. *Creat Nurs*. 2017;23(3):184-191.
62. Murphy M, Curtis K, McCloughen A. What is the impact of multidisciplinary team simulation training on team performance and efficiency of patient care? An integrative review. *Australas Emerg Nurs J*. 2016;19(1):44-53.
63. Patel A, Nouraei SA. Transnasal humidified rapid-insufflation ventilator exchange (THRIVE): a physiological method of increasing apnoea time in patients with difficult airways. *Anaesthesia*. 2015;70(3):323-329.
64. Jaber S, Monnin M, Girard M, et al. Apnoeic oxygenation via high-flow nasal cannula oxygen combined with non-invasive ventilation preoxygenation for intubation in hypoxaemic patients in the intensive care unit: the single-centre, blinded, randomised controlled OPTINIV trial. *Intensive Care Med*. 2016;42(12):1877-1887.
65. Neligan PJ, Malhotra G, Fraser M, et al. Continuous positive airway pressure via the Boussignac system immediately after extubation improves lung function in morbidly obese patients with obstructive sleep apnea undergoing laparoscopic bariatric surgery. *Anesthesiology*. 2009;110(4):878-884.
66. Zoremba M, Kalmus G, Begemann D, et al. Short term non-invasive ventilation post-surgery improves arterial blood-gases in obese subjects compared to supplemental oxygen delivery—a randomized controlled trial. *BMC Anesthesiol*. 2011;11:10.
67. Patient Safety Movement. Actionable patient safety solutions. Challenge 8: Airway Safety. <https://patientsafetymovement.org/?actionable-solutions/?challenge-solutions/?airway-safety/> (<https://patientsafetymovement.org/actionable-solutions/challenge-solutions/airway-safety/>). Accessed April 10, 2019.

Copyright © 2019 McMahon Publishing, 545 West 45th Street, New York, NY 10036. Printed in the USA. All rights reserved, including the right of reproduction, in whole or in part, in any form.




(/download/Unplanned_ANSE0819_WM.pdf)

Download to read this article in PDF document:

 [Unplanned or Accidental Extubation In the Perioperative Environment \(/download/Unplanned_ANSE0819_WM.pdf\)](/download/Unplanned_ANSE0819_WM.pdf)

This article is in PDF format and it requires Adobe Reader. If you do not have Adobe Reader installed on your computer then please download and install from the link below.

 [Download Adobe Reader \(http://get.adobe.com/reader/\)](http://get.adobe.com/reader/)

Related Keywords

[airway management \(/Tag/airway-management/399\)](/Tag/airway-management/399)

[extubation \(/Tag/extubation/1307\)](/Tag/extubation/1307)

- HIDE COMMENTS

B **U** **I**

Disclaimer: This board is intended for non-commercial reader comments. The posting of advertisements or solicitation of any kind is prohibited. Contributors are expected to conduct themselves in an appropriate manner, refraining from personal attacks and profane, coarse or abusive language. Hit the "Report Abuse" button to report comments that violate the rules and spirit of this board. www.anesthesiologynews.com reserves the right to remove without any notice comments that violate the intentions of this board. It is not intended to provide specific medical advice or to take the place of written law or regulations. The owners of www.anesthesiologynews.com assume no liability for the comments made on this board.

SUBMIT

More in Review Articles (/Section/Review-Articles/161)

Rural Airway Management: A Case of Severe Angioedema in a Small Community Hospital (/Review-Articles/Article/08-19/Rural-Airway-Management-A-Case-of-Severe-Angioedema-in-a-Small-Community-Hospital/55654)

We present a case of severe angioedema in a small community.....
AUGUST 28, 2019

Shocking Tales of a Horribly Soiled Airway (/Review-Articles/Article/08-19/Shocking-Tales-of-a-Horribly-Soiled-Airway/55659)

James Ducanto, MD, and Chris Root, NRP, show how the SALAD.....
AUGUST 26, 2019

Airway Management In Highly Infectious Diseases: The Reemergence of Measles (/Review-Articles/Article/08-19/Airway-Management-In-Highly-Infectious-Diseases-The-Reemergence-of-Measles/55660)

Measles serves as an excellent, albeit unfortunate, example of a highly.....
AUGUST 23, 2019

MORE (/Section/Review-Articles/161)

Most Read

Looming Anesthesiologist Shortage Fuels High Market Demand (/PRN-/Article/08-19/Looming-Anesthesiologist-Shortage-Fuels-High-Market-Demand/55607)

AUGUST 13, 2019 PRN (/Section/PRN-/21)

Best Neuraxial Technique for Hip and Knee Arthroplasty Is Spinal Anesthesia (/Pain-Medicine/Article/07-19/Best-Neuraxial-Technique-for-Hip-and-Knee-Arthroplasty-Is-Spinal-Anesthesia/55293)

JULY 16, 2019 PAIN MEDICINE (/Section/Pain-Medicine/2)

New Airway Keeps Jaw Protruded, Tongue in Place (/Technology/Article/08-19/New-Airway-Keeps-Jaw-Protruded-Tongue-in-Place/55614)

AUGUST 6, 2019 TECHNOLOGY (/Section/Technology/8)

Profound Lip and Tongue Swelling After Anesthesia Induction (/Clinical-Anesthesiology/Article/07-19/Profound-Lip-and-Tongue-Swelling-After-Anesthesia-Induction/55294)

JULY 10, 2019 CLINICAL ANESTHESIOLOGY (/Section/Clinical-Anesthesiology/1)

Tweets by @anesthesianews



Anesthesiology News
@anesthesianews

Collaborative announced focused on increasing awareness of the magnitude of the problem of unplanned [#extubation](#) [@AmerAcadPeds](#) [@AANAwebupdates](#) [@ACEPNow](#) [@society_asa](#) [@socforairway](#) [@APSPforg](#) [@PediAnesthesia](#) [@SCCM](#)
anesthesiologynews.com/Review-Article...

Unplanned or Accidental Extubation in the Perioperative ...
More than 120,000 incidents of unplanned or uncontrolled ext...
anesthesiologynews.com

[Embed](#)

[View on Twitter](#)

Monographs & Whitepapers (/Section/Monographs-and-Whitepapers/443)

The Roles of the Anesthesiologist, Pharmacist, and Nurse in Detecting and... (Monographs-and-Whitepapers/Article/06-19/The-Roles-of-the-Anesthesiologist-Pharmacist-and-Nurse-in-Detecting-and-Preventing-Opioid-Diversion-and-Misuse/55237)

Diversion of controlled substances from the health care setting remains a significant problem. Anesthesiologists, pharmacists, and nurses...



Clinical and Economic Impact of Postoperative Nausea and Vomiting (/Monographs-and-Whitepapers/Article/03-19/Clinical-and-Economic-Impact-of-Postoperative-Nausea-and-Vomiting/54418)

This monograph describes the epidemiology, pathophysiology, and burden of postoperative nausea and vomiting, and discusses current strategies...

Archives (/Archive)

[July 2019 \(/Archive/July-2019\)](#)

[June 2019 \(/Archive/June-2019\)](#)

[May 2019 \(/Archive/May-2019\)](#)

[April 2019 \(/Archive/April-2019\)](#)

[MORE \(/Archive\)](#)



<https://www.AnesthesiologyNews.com>

Anesthesiology News, now in its 45th year, has been the best-read publication for the specialty for the last 22 years. All 51,911 anesthesiologists and anesthesiology residents in the United States receive a copy every month!

HOME

<https://www.AnesthesiologyNews.com/LandingPage>

[ABOUT https://www.AnesthesiologyNews.com/About](https://www.AnesthesiologyNews.com/About)

SUBSCRIPTION

ADVERTISE

<https://www.AnesthesiologyNews.com/Advertise>

CONTACT

<https://www.AnesthesiologyNews.com/Contact>

ARCHIVES

<https://www.AnesthesiologyNews.com/Archive>

[RSS FEED \(/RSS/RSS.xml\)](#)

PRIVACY POLICY

<https://www.anesthesiologynews.com/privacypolicy>

eNewsletter

[Click here](#)

<https://www.AnesthesiologyNews.com/eNewsSignup> (<http://www.anesthesiologynews.com>)

to sign up to receive our eNewsletter.

Social Media

[anesthesiologynews](#)

<https://www.facebook.com/anesthesiologynews>

[@anesthesianews](#)

<https://twitter.com/anesthesianews>

[anesthesiologynews](#)

<https://www.linkedin.com/company/anesthesiologynews/>

McMahon Websites



Anesthesiology News



<http://www.anesthesiologynews.com>



Clinical Oncology News



<http://www.clinicaloncology.com>



Gastroenterology & Endoscopy News



<http://www.Gastroendonews.com>



General Surgery News



<http://www.Generalsurgerynews.com>



Infectious Disease Special Edition



<http://www.idse.net>



Pain Medicine News



<http://www.painmedicineneeds.com>



Pharmacy Practice News



<http://www.Pharmacypracticeneeds.com>



Specialty Pharmacy Continuum



<http://www.Specialtypharmacycontinuum.com>



CMEZone.com (<http://www.CMEZone.com>)



McMahon Custom Media



<http://custommedia.mcmahonmed.com/>



McMahon Group



<http://www.mcmahonmed.com/>



Pharmacy Technology Report



<http://www.pharmacytechnologyreport.com/>



OR Management News



<https://www.ormanagement.net/>

My Account

LOGOUT

<https://www.AnesthesiologyNews.com/Layout/Logout>

MANAGE MY ACCOUNT

<https://www.AnesthesiologyNews.com/manage-account>

Copyright © 2004-2019 McMahon Publishing unless otherwise noted.

Anesthesiology News and AnesthesiologyNews.com are part of McMahon Publishing.

All rights reserved. Reproduction in whole or in part without permission is prohibited.